



---

## COMMON SKINK *Eutropis carinata* (REPTILIA: SCINCIDAE) FEED ON ENDEMIC SEMI-SLUG *Ratnadvipia irradians* (LIMACOIDEA: ARIOPHANTIDAE)

---

Sectional Editor: Sujan Henkanaththegedara

Submitted: 30 August 2009, Accepted: 12 October 2009

D. M. S. Suranjan Karunarathna<sup>1</sup> and A. A. Thasun Amarasinghe<sup>2</sup>

<sup>1</sup>IUCN Sri Lanka Country office, No. 53, Horton place, Colombo 07, Sri Lanka

<sup>2</sup>Taprobanica Nature Conservation Society, 146, Kendalanda, Homagama, Sri Lanka

<sup>2</sup>Corresponding author: aathasun@gmail.com

The skinks belong to the genus *Eutropis* is widely distributed in South and South East Asia and is represented in Sri Lanka by seven species, the largest and the most common of which is common skink *Eutropis carinata* (Schneider, 1801) (Das *et al.*, 2008). In Sri Lanka *E. carinata* is widely distributed in open areas, closed canopy forest, home gardens and plantations in wet and dry zones below 1,000 m altitude (Das & de Silva, 2005). The Sri Lankan endemic semi-slug genus *Ratnadvipia* consists of two species, *Ratnadvipia irradians* (Pfeiffer, 1853) and *Ratnadvipia karui* Raheem & Naggs, 2006. *Ratnadvipia* is almost exclusively confined to the tropical lowland rain forests, intermediate zone and up to the suitable habitats within the dry zone (Raheem & Naggs, 2006). Although snails are usually not an item of skink diet, we have observed a semi-slug in *E. carinata* gut.

On 30<sup>th</sup> December 2008, a dry sunny day, first author found a dead (road kill) of *Eutropis carinata* adult male (snout-vent length 143 mm, tail length 202 mm) on the rocky road at Morningside area, Ratnapura District (06°24'N, 80°38'E, altitude 1030 m) at 07:15 hr (temperature 23 °C, humidity 69%). The specimen was fresh with severe head damage. The abdomen of the skink was dissected and investigated the gut contents. The lumen of the stomach consisted of an undigested adult specimen of *Ratnadvipia irradians* (total length of the body: about 60 mm, maximum width of the shell: about 18 mm) (Fig. 1) in addition to six half digested Coleopterans.

We have frequently observed *R. irradians* at Morningside Forest. This species is mostly observed on the leaves of cardamom (*Elettaria cardamonium*) bushes, on shrubs as well as on rotting

logs at night (normally 0.5-2 m high) also during the dry season under leaf litter, under rocks and inside rotting logs (Bahir and Gabadage Pers. com. Feb. 2009). *E. carinata* is commonly found under leaf litter and observed on rock surfaces, but we have not previously observed this skink feeding on *R. irradians* or on any other land snails. Additionally we observed a live *R. irradians* on the scats of a Golden Jackal, *Canis aureus lanka* on the same road (Fig. 2). The scats were wet and with a fungus growth. Although we could not verify that *Ratnadvipia* feeding on jackal scat, it may have been feeding on the fungus.




Fig. 01: undigested adult specimen of *R. irradians*



Fig. 02: *R. irradians* on the scats of Jackal

This is the first observation of *E. carinata* feeding on a land snail *R. irradians*. Further investigation is needed to establish if *R. irradians* is a regular item of the diet of *E. carinata* and to clarify what *R. irradians* feeds when on Jackal scats. Data on the diet and feeding habits of Sri Lankan and South Asian land snails or skinks are sparse and further observations are needed to verify their feeding habits. It is interesting to examine food and feeding habits of *E. carinata* on its range especially in

mainland India. Since *Ratnadvipia* is endemic to Sri Lanka, it is worthwhile to study the feeding interactions between this endemic semi-slug and other skinks in general including *E. carinata*.

#### Acknowledgements

We are grateful to Fred Naggs, Dinarzarde Raheem, Indraneil Das, M. M. M. Najim and M. M. Bahir for reviewing the manuscript and providing valuable comments. First author thanks Mendis Wickramasinghe for his support. Finally we thank Dinesh E. Gabadage, Umanga Gunasekara, Sampath Kolambage, Udaya Chanaka, Amila Chanaka, Dulan Ranga and Saman Nawaratne for their support.

#### Literature Cited

Das, I. and A. de Silva, 2005. *Photographic Guide to Snakes and other Reptiles of Sri Lanka*. New Holland Publishers (UK) Ltd., London: 144.

Das, I., A. de Silva, and C. C. Austin, 2008. A new species of *Eutropis* (Squamata: Scincidae) from Sri Lanka. *Zootaxa*, 1700: 35–52.

Pfeiffer, L., 1853. *Monographia Heliceorum Viventium*. Volume 3.

Raheem, D. and F. Naggs, 2006. The Sri Lankan endemic semi-slug *Ratnadvipia* (Limacoidea: Ariophantidae) and a new species from southwestern Sri Lanka. *Systematics and Biodiversity*, 4 (1): 99–126.

Schneider, J. G., 1801. *Historiae Amphibiorum naturalis et literariae. Fasciculus secundus continens Crocodilos, Scincos, Chamaesauras, Boas, Pseudoboas, Elapes, Angues, Amphisbaenas et Caecilias*. Frommani, Jena: 374.